

IX INTERNATIONAL SYMPOSIUM ON AGRICULTURAL SCIENCES

24th September 2020
Banja Luka
Bosnia and Herzegovina

BOOK OF ABSTRACTS



AGRORES 2020



BOOK OF ABSTRACTS

IX International Symposium on Agricultural Sciences "AgroReS 2020"
24 September, 2020; Banja Luka, Bosnia and Herzegovina

Publisher

University of Banja Luka
Faculty of Agriculture
University City
Bulevar vojvode Petra Bojovića 1A
78000 Banja Luka, Republic of Srpska, B&H

Editor in Chief

Željko Vaško

Technical Editors

Biljana Rogić

Circulation

online on the website <https://agrores.net/zbornici/>

CIP - Каталогизacija у публикацији

Народна и универзитетска библиотека

Републике Српске, Бања Лука

631(048.3)

INTERNATIONAL Symposium on Agricultural
Sciences (9 ; Banja Luka ; 2020)

Book of Abstracts [Elektronski izvor] / 9th
International Symposium on Agricultural Sciences
"AgroReS 2020", 24 September, 2020, Banja Luka,
Bosnia and Herzegovina ; [organizer University of
Banjaluka, Faculty of Agriculture ; editor in chief Željko
Vaško]. - Banja Luka : Faculty of Agriculture =
Poljoprivredni fakultet, 2020

Način pristupa (URL): <https://agrores.net/zbornici/>

ISBN 978-99938-93-63-9

COBISS.RS-ID 129416961

02_20

Bionomy of *Parthenolecanium persicae* (Fabricius) (Hemiptera: Coccidae) on grapevine in Serbia

Marina Dervišević¹, Aleksandar Stojanović², Lazar Sivčev³, Draga Graora⁴

¹*Biounik d.o.o., Research and Development Center, Belgrade, Republic of Serbia*

²*Natural History Museum, Belgrade, Republic of Serbia*

³*Institute for plant protection and environment, Belgrade, Republic of Serbia*

⁴*University in Belgrade - Faculty of Agriculture, Belgrade, Republic of Serbia*

Corresponding author: Marina Dervišević, marina.dervisevic@agrounik.rs

Abstract

Parthenolecanium persicae (Fabricius) (Hemiptera: Coccidae) is a palearctic species found on over 70 host plants, with the highest damage being recorded on grapevine and plums. In Serbia, this one is more abundant on extensive orchards or physiologically weakened plants. Bionomy of *P. persicae* was studied from 2015 to 2019 in three localities: Aleksandrovac, Belgrade and Neštin. Sampling of plant material was performed throughout the year, in interval of 7 to 10 days during vegetation, or, once in month during plant dormancy. From each infested plant, five one- or two- years old twigs 20 cm long were taken. Sampled plant material with scale insect colonies was brought to the laboratory, where it was examined for scale insects presence and then deposited in glass cylinders for the rearing of scales insects and natural enemies. The determination of the eclosed parasitoids was performed by eng. Aleksandar Stojanović. *P. persicae* developed one generation per year and overwintered as second instar larva on grapevine shoot and trunk. Females emerged during April and early May, and oviposition was recorded in mid-May. Each female layed on average $2001 \pm 3,4$ white eggs. First instar larvae appeared during June. In the summer, they feed mainly on young shoots, leaves and leaf petioles. In late August and early September, second instar larvae were formed, which continue to feed until October when they migrated to shoots or hidden places on the trunks where they overwintered. Numerous colonies of *P. persicae* was recorded in the Belgrade locality, which resulting in discoloration, drying and premature leaf fall. In the vineyards of Aleksandrovac and Neštin, scale insects was present on individual plants, with no visible symptoms of damage. During our research, on the *P. persicae* we found 7 species of natural enemies. Of the parasitoids (Hymenoptera: Chalcidoidea), 6 species were eclosed: *Coccophagus lycimnia* (Walker), *Eunotus obscurus* Masi, *Marietta picta* (Andre) (Aphelinidae), *Metaphycus dispar* (Mercet), *Metaphycus insidiosus* (Mercet), *Metaphycus maculipennis* (Timberlake),

(Encyrtidae), and from predators, the species *Exochomus quadripustulatus* L. (Coleoptera: Coccinellidae).

Key words: Bionomy, *Parthenolecanium persicae*, grapevine, Serbia